
VISION

PRE-EXISTING CONDITIONS

Any pre-existing condition, especially a vestibular condition, should be discussed with the doctor prior to the operation.

ARTICLE

52

**DID THIS ARTICLE
HELP YOU?
SUPPORT VEDA @
VESTIBULAR.ORG**

5018 NE 15th Ave.
Portland, OR 97211
1-800-837-8428
info@vestibular.org
vestibular.org

Post-Operative Balance Issues Following Cataract Surgery

By Nathan Davis, OD, COVD, NORA

The visual and vestibular systems work together to allow a person to move about. Because these two systems are so closely integrated, any disruption in the processing of either system can have a negative impact on a person's sense of balance.

Cataract surgery can cause a dramatic change in the way a person visually experiences their world, and most of the time the effect is positive. However, if the brain is used to seeing the world from a certain perspective, and that perspective has suddenly been shifted, the disruption in the visual system can in turn cause sensory incoherence with the visual-vestibular connection.

PATIENTS WHO HAVE A PRE-EXISTING VESTIBULAR DYSFUNCTION ARE MORE AT RISK AFTER CATARACT SURGERY.

There are also various visual characteristics that may increase the likelihood of disrupting the visual-vestibular connection. Patients with significant corrections in their eyeglass prescription may have trouble adjusting to a new prescription, especially during the time between the first and second cataract surgery. Between the two procedures the prescription in one eye will typically be neutralized while the other eye remains the same. The difference between the two prescriptions can be hard for patients to adjust to at first. Any feelings of dizziness or nausea are typically resolved with the surgery for the second eye, but in rare cases the disruption can cause symptoms that may be hard to overcome. The real problem arises when the cataracts don't mature symmetrically, and one eye needs surgery well before the other eye.

OUTCOMES

Due to the technological advances of the implantable lens that is used during cataract surgery, a patient now has many different options concerning the visual outcome they desire. More specifically, there are options available today that allow a patient to see in the distance and up close without the aid of glasses, at least to some degree.



COMMON OUTCOMES OF CATARACT SURGERY

Monovision

Purposely sets one eye to focus on distance objects, while the other eye focuses on near targets.

Multifocal intraocular

Allows the patient to use both eyes to focus on objects up close and far away.

Distance only

Have both implants focused for distance, and to use glasses to see up close.

The two most popular options are multifocal intraocular lenses and a monovision outcome. Multifocal intraocular lenses allow the patient to use both eyes to focus on objects up close and far away. Monovision purposely sets one eye to focus on distance objects, while the other eye focuses on near targets. Both visual experiences are far different (and less efficient) than normal human vision, and as such can be difficult for the brain to adjust to, especially for a patient with a vestibular disorder.



KEY TO SUCCESS: COMMUNICATION

The key to success centers on communication between the doctor and the patient before surgery. **Any pre-existing condition, especially a vestibular condition, should be thoroughly explained** with all options explored. For most patients the safest result would be to choose to have both implants focused for distance, and to use glasses to see up close. This is generally the safest option as it most closely mimics how our vision is naturally used, emphasizing symmetry between the two eyes and encouraging our brain to use both eyes together. This set-up should also provide the smallest amount of disruption to the visual-vestibular connection. Extra caution should be given to the patient who is considering a multifocal implant or a monovision result, as well as the patient who is only having one eye operated on for the foreseeable future. Contact lens simulations before the surgery can be used to mimic the multifocal and monovision outcomes, and should be considered, if possible.

©2017 Vestibular Disorders Association
VeDA's publications are protected under copyright.
For more information, see our permissions guide at
vestibular.org. ***This document is not intended as a
substitute for professional health care.***



[illegible]

5018 NE 15th Ave. Portland, OR 97211
1-800-837-8428 info@vestibular.org vestibular.org

You can ensure that educational articles like this continue to be available to vestibular patients like you by making a tax-deductible gift to VeDA today.

One-time gift: ☐ \$40 ☐ \$50 ☐ \$75 ☐ \$100 ☐ \$250 ☐ other

Monthly gift: ☐ \$10 ☐ \$15 ☐ \$25 ☐ \$35 ☐ \$50 ☐ other

☐ Check this box if you prefer that your donation remain anonymous.

Donations gladly accepted online at <http://vestibular.org>. Check or money order in US funds, payable to VeDA.

Visa	MC	Amex	Discover	<div><div></div><div>Card number</div></div>	<div><div></div><div>Exp. date</div></div>	<div><div></div><div>CVV code</div></div>
------	----	------	----------	--	--	---

Billing address of card (if different from mailing information)

Name _____ Telephone _____ Email _____
Address _____ City _____ State/Province _____ Zip _____
Country _____