



VESTIBULAR

DISORDERS ASSOCIATION

PO Box 13305 · PORTLAND, OR 97213 · FAX: (503) 229-8064 · (800) 837-8428 · INFO@VESTIBULAR.ORG · WWW.VESTIBULAR.ORG

Vestibular Injury

Compensation, De-compensation, and Failure to Compensate

By Thomas E. Boismier, MPH, The Balance Care Center, Fort Wayne, Indiana with the Vestibular Disorders Association

The vestibular system includes the inner ear balance organs and the parts of the brain that coordinate and process balance information. The balance organs contribute vital sensory information about motion, equilibrium, and spatial orientation. In addition, the vestibular system helps maintain clear vision during movements via an automated function called the *vestibulo-ocular reflex* (VOR).

The vestibular system can be compromised in many ways. Viral infections (labyrinthitis and vestibular neuritis), disorders that affect inner ear fluid levels (Ménière's disease and secondary endolymphatic hydrops), trauma from head injury, benign tumors (acoustic neuromas), and age-related degeneration can all cause permanent damage to it.

The body has limited ability to repair damage to the vestibular organs, although the body can often recover from vestibular injury by having the part of the brain that controls balance recalibrate itself to compensate. This compensation process occurs naturally in most people; however, when the compensation process is not successful, vestibular rehabilitation therapy may be required.

Acute (immediate) compensation

When a sudden injury occurs to the balance organs on one side, the patient may feel very sick for hours to a few days with vertigo (a spinning feeling), lightheadedness (dizziness), and unsteadiness, often accompanied by sweating, nausea, and vomiting. This is because the spatial orientation signals being sent to the brain from both ears are no longer equal and opposite, and the brain interprets the difference as constant movement.

Researchers theorize that after this initial period, the brain recognizes that the signals being received from the ears are incorrect and turns the signals off through a process called the *cerebellar clamp*. When the clamp is in place, the vertigo and much of the sick feelings improve. However, the patient feels unsteady while standing, because the vestibular signals normally used to maintain balance have been turned off.

The patient may also report dizziness or blurred vision with head movements because the VOR no longer functions properly. Information provided by proprioception (the sense of pressure at the bottom of the feet) and vision is

part of the natural process of maintaining balance. So without vestibular input, the patient can still walk but will feel unsteady and may fall in the dark or on soft or bumpy surfaces such as a thickly carpeted floor, grass lawn, or gravel path.

At this point, most patients are well enough to get out of bed and visit a doctor. The doctor sees a person who is not experiencing dizziness or vertigo but whose gait is *ataxic*, meaning awkward and uncoordinated. If the patient is not given an opportunity to clearly describe what has happened, he or she may be immediately referred to a neurologist to rule out stroke because of this ataxic gait.

The cerebellar clamp may persist for days to a few weeks after the initial injury. If balance testing is performed during this phase, test results may incorrectly suggest that the patient has damage to both sides of the balance system, because the cerebellar clamp reduces the eye movements (VOR) that are typically evaluated during balance testing.

Chronic (long-term) compensation

During the acute compensation phase, the cerebellum slowly releases the clamp, gradually allowing more signals from the inner ear to pass to the brain. The brain can then fine-tune its calculations in order to account for differences between the signals received from damaged versus undamaged vestibular organs. Such compensation can't occur if the cerebellar clamp remains in place.

For most patients, the movements made during normal daily activities are enough to achieve chronic (long-term) compensation, usually in two to four weeks after the injury has occurred. Once this chronic compensation process is complete, the patient is essentially symptom-free. If unsteadiness and/or motion-provoked dizziness persist after that time, compensation is not complete and the physician may prescribe a program of vestibular rehabilitation therapy (VRT).

VRT is a treatment program administered by a specially-trained physical or occupational therapist. It is designed to (1) desensitize the balance system to problematic movements and (2) enhance the fine-tuning required for long-term compensation. This is accomplished with exercises that involve small, controlled, and repeated "doses" of the movements and activities that provoke dizziness. VRT is most effective when administered by a therapist who has specific training and experience in this specialized form of therapy.

De-compensation

It's important to remember that even after symptoms go away, the balance system remains injured; the brain has simply adapted to the injury. For many patients, dizziness and vertigo will return months or years after compensating for a balance system injury. It is critical for the physician to find out what type of dizziness and vertigo the patient has.

If the patient describes another severe attack of spinning with unsteadiness and nausea lasting hours to days, this suggests that a second injury has occurred to the balance system, such as another viral infection or an attack of Ménière's disease or secondary endolymphatic hydrops. These conditions require diagnosis and medical treatment. If the patient reports that dizziness occurs after particular movements and lasts seconds to a few minutes, this suggests that de-compensation has occurred. *De-compensation* simply means that the brain has 'forgotten' the fine-tuning it developed during the chronic compensation phase described above.

Events that can provoke de-compensation include a bad cold or the flu, minor surgery, or anything that interrupts normal daily activity for a few days, such as long vacations. Recovery after de-compensation is exactly like the recovery that occurs during the chronic compensation phase. Movements and activities are the stimuli the brain needs to fine-tune the system. In our balance center, we routinely counsel patients to keep their VRT exercise program instructions in a drawer even after they recover so that they can begin the exercises immediately if symptoms return. Usually recovery after de-compensation is quicker than the time it takes to recover after the initial injury to the balance system.

Failure to compensate

Two things are required in order to compensate for an injury. First, the brain must receive signals from the balance organs. Thus, movements must not be avoided, because movements create the signals the brain needs to compensate for the injury. Second, the balance areas of the brain must be capable of change.

During the early stages of dizziness, many physicians counsel their patients to avoid quick movements and reduce their activities. Most patients will be prescribed one or more anti-dizziness medications such as Antivert (meclizine), Valium (diazepam), Xanax (alprazolam), Phenergan (promethazine), or Compazine (prochlorperazin). This is fine during the acute stages of a dizziness problem in order to reduce symptoms that persist for hours or days even when the patient is not moving. However, once the acute phase is past, inactivity and medications can interfere with the long-term compensation process.

Any medication that makes the brain sleepy, including all of the anti-dizziness medications, can slow down or stop the process of compensation, so they are often not appropriate for long-term use. Most patients who fail to compensate are found to either be strictly avoiding certain movements, using anti-dizziness medications daily, or both. Treatment includes VRT and gradual reduction and eventual elimination of these medications.

Brain damage caused by stroke, head injury, or other causes can slow down or stop the natural compensation process. It is difficult to predict which patients with brain injury will improve or to what degree, so all patients should be given the chance to improve through a VRT

Further reading

Another helpful document available from the Vestibular Disorders Association is: Vestibular Rehabilitation – An Effective Evidence-Based Treatment.

© 2009 Vestibular Disorders Association

program. In our balance center we use several different measures of symptoms and functional capabilities in order to assess progress repeatedly as treatment goes on. As long as a patient continues to show improvement, even if it is gradual, treatment should be continued.

VEDA's publications are protected under copyright. For more information, see our permissions guide at www.vestibular.org.

This document is not intended as a substitute for professional health care.



VESTIBULAR DISORDERS ASSOCIATION

PO Box 13305 · PORTLAND, OR 97213 · FAX: (503) 229-8064 · (800) 837-8428 · INFO@VESTIBULAR.ORG · WWW.VESTIBULAR.ORG

Did this free publication from VEDA help you?

Thanks to VEDA, vestibular disorders are becoming recognized for their impacts on lives and our economy. We see new diagnostic tools and research studies, more accessible treatments, and a growing respect for how life-changing vestibular disorders can be.

VEDA provides tools to help people have a better quality of life: educational materials, support networks, professional resources, and elevated public awareness.

Your support of VEDA matters. Please help us to continue providing such great help by becoming a member or donor.

Members receive an information packet; discounts on purchases; a subscription to VEDA's newsletter, *On the Level*, containing information on diagnosis, treatment, research, and coping strategies; and the option of communicating directly with others who understand the personal impacts of a vestibular disorder. Professional members also receive the option to list training opportunities on our site, bulk-discounted prices on patient education materials, and a listing on VEDA's provider directory, the only of its kind serving patients seeking help from a vestibular specialist.

SUPPORT VEDA

Membership, 1-year

\$ 40 ... Basic

\$110 ... Professional

Memberships include electronic & online newsletter & free publications. For hard copies, include optional shipping fees.

\$ 5 ... Shipping (domestic)

\$ 15 ... Shipping (international)

\$ _____ Please indicate your desired subscription amount here.

Optional Contribution

I'd to support VEDA with a donation (instead of or in addition to membership).

\$ _____ Please indicate your desired subscription amount here.

Check this box if you prefer that your donation remain anonymous.

\$

Total

PAYMENT INFORMATION

If you prefer, you can make your purchases online at <http://www.vestibular.org>.

Check or money order in US funds, payable to VEDA (enclosed)

Visa

MC

Amex

_____ Card number

_____ Exp. date (mo./yr.)

_____ Billing address of card (if different from mailing information)

MAILING INFORMATION

Name _____

Address _____ City _____

State/Province _____ Zip/Postal code _____ Country _____

Telephone _____ E-mail _____