
DIAGNOSIS

TESTING

Diagnostic testing is an essential step to uncover the root source of vestibular problems.

ARTICLE

026

**DID THIS ARTICLE
HELP YOU?
SUPPORT VEDA @
VESTIBULAR.ORG**

5018 NE 15th Ave.
Portland, OR 97211
1-800-837-8428
info@vestibular.org
vestibular.org

Diagnostic Tests for Vestibular Disorders

By the Vestibular Disorders Association with Kelsey Hatton, Au.D., CCC-A, Kamran Barin and Daniel J. Romero, AuD, PhD

VESTIBULAR SYSTEM

Separate from your organ of hearing, your inner ear is also home to structures that help with balance, called the vestibular system, consisting of five organs in each ear. These organs and the associated nerves and brain centers form a complex system that serves many functions through connections to the brain, eyes, and feet. Thus, a thorough evaluation of the inner ear may require several different kinds of tests.

The vestibular and visual systems are connected to each other and to the muscles in the eyes and neck that help maintain good balance. Head movements or other stimulation of the inner ear sends signals through the nervous system to control eye muscle movements. This forms a reflex pathway called the vestibulo-ocular reflex, or VOR. This system is designed to generate eye movements that maintain clear vision when the head is in motion.

HOW ARE VESTIBULAR DISORDERS DIAGNOSED?

Many vestibular tests used when diagnosing vestibular disorders use equipment to monitor the eyes for normal and abnormal movements when the vestibular system is actively being stimulated.

Diagnosing vestibular disorders is challenging, for many reasons. However, doctors use information from a person's medical history and findings from a physical examination as a basis for ordering diagnostic tests to assess the vestibular system function and to rule out alternative causes of symptoms. Most people tolerate these tests well. However, sometimes the tests are fatiguing and can result in temporary unsteadiness. Vestibular testing does not affect your condition or



VNG Binocular Goggles



cause future vestibular episodes.

TESTS FOR DIAGNOSING VESTIBULAR DISORDERS

ELECTRO/VIDEO-NYSTAGMOGRAPHY (ENG OR VNG)

Electronystagmography (ENG) refers to a group of tests or test battery, and uses small electrodes placed over the skin around the eyes during testing. Videonystagmography (VNG) refers to the same test battery, but uses a set of video goggles to monitor the eyes. Both video cameras (VNG) and the electrodes (ENG) can measure eye movements to evaluate signs of vestibular dysfunction or neurological problems and are a key step in diagnosing vestibular disorders. Generally these tests are performed in a room



Caloric Air Fx Testing

that is dark or with low lighting. The examiner asks random questions that are meant to distract the person being tested and keep them alert. ENG/VNG tests are the most common set of tests administered to people with dizziness, vertigo, and/ or imbalance.

Parts of the ENG/VNG test battery evaluate the movement of the eyes as they follow different visual targets, which may be projected on either a light bar or TV screen. Other parts of the ENG/VNG observe eye movements as the head is placed in different positions. A third component of the ENG/VNG is called the caloric test, which uses water or air to change the temperature within the ear canal to stimulate part of the inner ear's vestibular organs. Air or water may be warmer or cooler than body temperature. In a healthy ear, this test does result in a sensation of motion and jerking eye movements (nystagmus) for a short

time. The caloric test evaluates the vestibular system at slow speeds.

ROTATIONAL TESTING

Rotational testing evaluates how well the eyes and inner ear work together at slow to moderate speeds. These tests are another key element to diagnosing vestibular disorders, and also use video goggles or electrodes to monitor eye movements. The head is rotated from side to side at slow to moderate speeds, and associated eye movements are analyzed. Like the ENG/VNG, rotation tests are performed in a room that is dark with the examiner asking random questions during testing. Rotational tests provide information beyond the ENG/VNG about how well the balance organs are functioning. Not all people in the diagnosis phase will require these tests.

There are different kinds of rotational tests: computerized rotary chair, auto head rotation, or a screening test. During computerized rotary chair tests, the patient sits in a motorized chair that swivels side to side at a controlled speed. In auto head rotation, the person being tested is asked to look at a fixed target and move his/her head back and forth or up and down for short periods of time. Screenings can be performed with the examiner watching the eyes while turning the subject side to side in a swivel chair.

VIDEO HEAD IMPULSE TESTING (vHIT)

vHIT evaluates how well the eyes and inner ears work together at speeds more consistent with natural everyday head movement. A small set of glasses with a camera are used to monitor eye movements. The vHIT test uses very small and quick movements of the head to evaluate reflex function, as opposed to the slow or moderate speeds used in rotation testing. Not all people in the diagnosis phase will require vHIT tests.

VESTIBULAR EVOKED MYOGENIC POTENTIAL (VEMP)

VEMP testing is used to evaluate whether certain vestibular organs that help us detect gravity (called the otolith organs) and their associated nerves are structurally intact. Responses in this test are measured from different muscles in the neck and around the eyes. VEMP testing uses adhesive, skin surface electrodes (like ENG or some rotational tests) and earphones (like those used during a hearing test). Sound is played





VEMP Test

for a few seconds through the earphones, which stimulates the otolith organs and activates muscle responses, and electrodes record the results.

COMPUTERIZED DYNAMIC POSTUROGRAPHY (CDP)



CDP Testing
Courtesy of Bertec

CDP tests how well someone maintains their balance under different environmental conditions. Maintenance of balance depends on sensory information from: the body's muscles/joints, eyes, and inner ears. This testing investigates relationships among these three sensory systems and records the balance and posture adjustments made when different scenarios are presented. This

test may also be used in a rehabilitative setting after a diagnosis has been determined and is not performed on all people in the diagnosis phase. CDP tests involve standing still on a platform. The platform may be still or able to shift, or a visual target may be still or able to move during testing. Pressure gauges under the platform record shifts in body weight (body sway) as the person being tested maintains balance under different conditions. A safety harness is worn as a precaution, should the patient lose their balance.

AUDIOMETRY (HEARING TESTS)

Audiometry measures hearing function. Hearing evaluations are an important part of vestibular diagnostics, because the inner ear contains both hearing and balance organs. More than one hearing test may be required when a person has a vestibular disorder, especially when there is evidence of hearing loss, a sensation of fullness in the ears, or tinnitus (ringing or noise in the ears).

The audiometric test battery is carried out in a sound-treated room. Earphones are used to present words and tones at different pitches and levels. A response is requested when these sounds are heard. Testing with words may include repeating words in a quiet room or when noise is playing.

Another part of a standard hearing test is tympanometry, which can help detect problems between the eardrum and the inner ear (a space known as the middle ear). Tympanometry uses a small earpiece that creates pressure and plays sound in the ear canal to gather information. The same equipment can also be used for acoustic-reflex testing, which measures the reflex of muscles in the middle ear in response to pressure and loud sound.

OTOACOUSTIC EMISSIONS (OAE)

OAE testing provides information about how the hair cells of the cochlea are working by measuring the responsiveness of hair cells to a series of clicks produced by a tiny speaker inserted into the ear canal. Most often this test is used to evaluate hearing for people who are unable to respond to a traditional hearing test (such as infants).

ELECTROCOCHLEOGRAPHY (ECoG)

ECoG measures a response to sound from the nervous system. It utilizes an earphone and



electrodes while the person being tested lays still in a comfortable position. Not all people in the diagnosis phase will need ECoG tests.

An earphone plays sound in the ear and an electrode measures a response. Different electrodes can be used in this test. Some may be adhesive, skin-surface electrodes. Others may fit in the ear canal like an earphone, while a third type of electrode is designed to gently rest against or touch the eardrum. A fourth type of electrode is a needle that is placed through the eardrum to touch the inner ear. Most clinics use the first three types of electrodes to measure an electrical signal while sound is playing.

AUDITORY BRAINSTEM RESPONSE TEST (ABR; OR BER, BSER, OR BAER)

The ABR measures how the nervous system responds to sound. The test setup and procedure is like the ECoG. Most often ABR is used to test hearing for people who are unable to respond for audiometry (such as infants). Occasionally this test is used when someone cannot have imaging performed (such as people with a metal plate in the body/brain).

Under certain circumstances, this test can indicate the presence of an acoustic neuroma (a rare, benign tumor of the vestibulo-cochlear nerve). It may also help identify conditions such as multiple sclerosis if they have affected the auditory pathway to the brain.

IMAGING

MAGNETIC RESONANCE IMAGING (MRI)

MRI uses a magnetic field and radio waves to produce cross-sectional images of body tissues being scanned. An MRI of the brain can reveal the presence of tumors, stroke damage, and other soft-tissue abnormalities that might cause dizziness or vertigo. MRIs of structures in and around the inner ear can be helpful in diagnosing vestibular disorders in some instances.

COMPUTERIZED AXIAL TOMOGRAPHY (CAT, OR CT)

A CT scan is an X-ray technique that is best for studying bony structures. The inner ear is inside of the skull's temporal bone on each side. These scans are often used to look for abnormalities around the inner ear, such as fractures or areas with thinning

bone.

OTHER TESTS FOR DIAGNOSING VESTIBULAR DISORDERS

Depending on your circumstances, other tests may be necessary to discover the cause of a balance disorder. Blood work, allergy tests, vision tests, and other exams may help rule out causes of imbalance that are unrelated to the vestibular system.

WHO PERFORMS VESTIBULAR TESTING?

There are many specialists who may be involved in diagnosing vestibular disorders. Generally, your primary care physician, ENT, neurologist, or neurologist will refer you to an audiologist for hearing or vestibular testing, a physical therapist for gait or balance related testing, or a radiologist for imaging testing. These specialists will send your test results back to your physician with a full report.

1. Campbell K. Essential Audiology for Physicians. San Diego: Singular Publishing Group; 1997.
2. Jacobson GP, Newman CW, Kartush JM. Handbook of Balance Function Testing. San Diego: Singular Publishing Group; 1993.
3. Jacobson GP, Shepard NT. Balance Function Assessment and Management. San Diego: Plural Publishing Inc; 2008. Second edition; 2014.
4. MacDougall HG, Weber KP, McGarvie LA, Halmagyi GM, Curthoys IS. The video head impulse test: diagnostic accuracy in peripheral vestibulopathy. *Neurology* 73 (14): 1134-1141. 2009.
5. McCaslin DL. Electronystagmography/ Videonystagmography (ENG/VNG). San Diego: Plural Publishing Inc; 2012.
6. Van den Hauwe L et al. Imaging in Patients with Vertigo. *JBR-BTR*. 1999;82:241-244.
7. Shepard NT, Telian SA. Practical Management of the Balance Disorder Patient. San Diego: Singular Publishing Group; 1996.

©2024 Vestibular Disorders Association
VeDA's publications are protected under copyright.
For more information, see our permissions guide at
vestibular.org. *This document is not intended as a
substitute for professional health care.*

